



HQExoTM **Exosome-A549**

Catalog: Exo-CH02

PRODUCT INFORMATION

Exosome derived from human non-small cell lung cancer cell line (A549 cell line) Exosomes are nanosized vesicles (30-160 nm) secreted by exocytosis by most cell types and contain specicargos, such as RNAs, lipids, and proteins. The cargos amount and composition of exosomes depend on the 1 type from which they are released, which making them useful for biomarker discovery and functional cherization. Exosomes have been isolated from cancer cell lines (human and mouse), which helps understand or growth microenvironments. Exosome derived from enormous model human cancer cell lines to improve studies of tumor growth and invasion signaling pathways as well as how these tumor exosomes function at an insight into antitumor research. HQExo TM standard exosomes could use as positive controls for exosometation and functional research, such as ELISA, FACS, WB. Lyophilization is useful for a long-term storated "C, and frozen liquid should be kept at -20°C to -80°C. Ultracentrifugation and precipitation techniques at a sinly used in exosome Isolation. It had been reported that both methods yielded extracellular vesicles in the erange of exosomes and included apoproteins, which can be used in downstream analyses. Nanoparticles king Analysis (NTA) is used for measuring exosome particles concentration, and WB or ELISA can be us exosomal biomarkers analysis. Creative Biostructure standard exosome products guarantee higher purity a uality to meet our customer research. Lyophilized powder/ frozen liquid. Reconstitute lyophilized exosome by adding deionized water for a desi inal concentration. Centrifuge before opening to ensure exosomes are at bottom, resuspend exosomes by p ing and/or vortex, please avoid bubbles. Centrifuge again and mix well for using. >1x10^8 particles Lyophilized powder store at 4 °C. Frozen liquid store at -20°C to -80°C. Recommended to avoid repeated		
Exosome derived from human non-small cell lung cancer cell line (A549 cell line) Exosomes are nanosized vesicles (30-160 nm) secreted by exocytosis by most cell types and contain speci cargos, such as RNAs, lipids, and proteins. The cargos amount and composition of exosomes depend on the ltype from which they are released, which making them useful for biomarker discovery and functional cheerization. Exosomes have been isolated from cancer cell lines (human and mouse), which helps understand or growth microenvironments. Exosome derived from enormous model human cancer cell lines to improve studies of tumor growth and invasion signaling pathways as well as how these tumor exosomes function at an insight into antitumor research. HQExo™ standard exosomes could use as positive controls for exosometation and functional research, such as ELISA, FACS, WB. Lyophilization is useful for a long-term storal 4°C, and frozen liquid should be kept at -20°C to -80°C. Ultracentrifugation and precipitation techniques at ainly used in exosome Isolation. It had been reported that both methods yielded extracellular vesicles in the erange of exosomes and included apoproteins, which can be used in downstream analyses. Nanoparticles king Analysis (NTA) is used for measuring exosome particles concentration, and WB or ELISA can be us exosomal biomarkers analysis. Creative Biostructure standard exosome products guarantee higher purity a uality to meet our customer research. Lyophilized powder/ frozen liquid. Reconstitute lyophilized exosome by adding deionized water for a desi inal concentration. Centrifuge before opening to ensure exosomes are at bottom, resuspend exosomes by p ing and/or vortex, please avoid bubbles. Centrifuge again and mix well for using. > 1x10^8 particles Lyophilized powder store at 4 °C. Frozen liquid store at -20°C to -80°C. Recommended to avoid repeated	Name	HQExo™ Exosome-A549
Exosomes are nanosized vesicles (30-160 nm) secreted by exocytosis by most cell types and contain specicargos, such as RNAs, lipids, and proteins. The cargos amount and composition of exosomes depend on the latype from which they are released, which making them useful for biomarker discovery and functional characterization. Exosomes have been isolated from cancer cell lines (human and mouse), which helps understant or growth microenvironments. Exosome derived from enormous model human cancer cell lines to improve studies of tumor growth and invasion signaling pathways as well as how these tumor exosomes function at an insight into antitumor research. HQExo TM standard exosomes could use as positive controls for exosometation and functional research, such as ELISA, FACS, WB. Lyophilization is useful for a long-term storate 4°C, and frozen liquid should be kept at -20°C to -80°C. Ultracentrifugation and precipitation techniques at an insight into antitumor research. It is a second an included apoproteins, which can be used in downstream analyses. Nanoparticles king Analysis (NTA) is used for measuring exosome particles concentration, and WB or ELISA can be use exosomal biomarkers analysis. Creative Biostructure standard exosome products guarantee higher purity at uality to meet our customer research. Lyophilized powder/ frozen liquid. Reconstitute lyophilized exosome by adding deionized water for a desitinal concentration. Centrifuge before opening to ensure exosomes are at bottom, resuspend exosomes by pring and/or vortex, please avoid bubbles. Centrifuge again and mix well for using. > 1x10^8 particles Lyophilized powder store at 4 °C. Frozen liquid store at -20°C to -80°C. Recommended to avoid repeated	Cat No.	Exo-CH02
cargos, such as RNAs, lipids, and proteins. The cargos amount and composition of exosomes depend on the latype from which they are released, which making them useful for biomarker discovery and functional characterization. Exosomes have been isolated from cancer cell lines (human and mouse), which helps understand or growth microenvironments. Exosome derived from enormous model human cancer cell lines to improve studies of tumor growth and invasion signaling pathways as well as how these tumor exosomes function at an insight into antitumor research. HQExo™ standard exosomes could use as positive controls for exosomediation and functional research, such as ELISA, FACS, WB. Lyophilization is useful for a long-term storal 4°C, and frozen liquid should be kept at -20°C to -80°C. Ultracentrifugation and precipitation techniques a ainly used in exosome Isolation. It had been reported that both methods yielded extracellular vesicles in the erange of exosomes and included apoproteins, which can be used in downstream analyses. Nanoparticles king Analysis (NTA) is used for measuring exosome particles concentration, and WB or ELISA can be us exosomal biomarkers analysis. Creative Biostructure standard exosome products guarantee higher purity a uality to meet our customer research. Lyophilized powder/ frozen liquid. Reconstitute lyophilized exosome by adding deionized water for a desirand inal concentration. Centrifuge before opening to ensure exosomes are at bottom, resuspend exosomes by ping and/or vortex, please avoid bubbles. Centrifuge again and mix well for using. > 1x10^8 particles	Source	Exosome derived from human non-small cell lung cancer cell line (A549 cell line)
inal concentration. Centrifuge before opening to ensure exosomes are at bottom, resuspend exosomes by p ing and/or vortex, please avoid bubbles. Centrifuge again and mix well for using. > 1x10^8 particles Lyophilized powder store at 4 °C. Frozen liquid store at -20°C to -80°C. Recommended to avoid repeated	Product Overview	Exosomes are nanosized vesicles (30-160 nm) secreted by exocytosis by most cell types and contain specifical cargos, such as RNAs, lipids, and proteins. The cargos amount and composition of exosomes depend on the cell type from which they are released, which making them useful for biomarker discovery and functional characterization. Exosomes have been isolated from cancer cell lines (human and mouse), which helps understand turn or growth microenvironments. Exosome derived from enormous model human cancer cell lines to improve the studies of tumor growth and invasion signaling pathways as well as how these tumor exosomes function and get an insight into antitumor research. HQExo TM standard exosomes could use as positive controls for exosome is olation and functional research, such as ELISA, FACS, WB. Lyophilization is useful for a long-term storage at 4°C, and frozen liquid should be kept at -20°C to -80°C. Ultracentrifugation and precipitation techniques are not ainly used in exosome Isolation. It had been reported that both methods yielded extracellular vesicles in the size of exosomes and included apoproteins, which can be used in downstream analyses. Nanoparticles Tracking Analysis (NTA) is used for measuring exosome particles concentration, and WB or ELISA can be used in exosomal biomarkers analysis. Creative Biostructure standard exosome products guarantee higher purity and cuality to meet our customer research.
Lyophilized powder store at 4 °C. Frozen liquid store at -20°C to -80°C. Recommended to avoid repeated	Form	Lyophilized powder/ frozen liquid. Reconstitute lyophilized exosome by adding deionized water for a desired inal concentration. Centrifuge before opening to ensure exosomes are at bottom, resuspend exosomes by pipet ing and/or vortex, please avoid bubbles. Centrifuge again and mix well for using.
Lyophilized powder store at 4 °C. Frozen liquid store at -20°C to -80°C. Recommended to avoid repeated	Concentration	> 1x10^8 particles
e-and-thaw cycles.	Storage	Lyophilized powder store at 4 °C. Frozen liquid store at -20°C to -80°C. Recommended to avoid repeated free e-and-thaw cycles.