



# HQExo™ Exosome-Goat Plasma exosome

## Catalog: Exo-EV-A-004

### PRODUCT INFORMATION

<b>Name</b>	HQExo™ Exosome-Goat Plasma exosome
<b>Cat No.</b>	Exo-EV-A-004
<b>Source</b>	Exosome derived from Goat Plasma
<b>Product Overview</b>	<p>Exosomes are nanosized vesicles (30-160 nm) secreted by exocytosis by most cell types and contain specific cargos, such as RNAs, lipids, and proteins. The cargo amount and composition of exosomes depend on the cell type from which they are released, which makes them useful for biomarker discovery and functional characterization. Exosomes have been isolated from cancer cell lines (human and mouse), which helps understand tumor or growth microenvironments. Exosome derived from animal plasma to improve the studies of veterinary diseases. These exosomes can be used to further diagnosis and therapeutics in veterinary pre-clinical and clinical studies. HQExo™ standard exosomes could be used as positive controls for exosome isolation and functional research, such as ELISA, FACS, WB. Lyophilization is useful for a long-term storage at 4°C, and frozen liquid should be kept at -20°C to -80°C. Ultracentrifugation and precipitation techniques are mainly used in exosome isolation. It has been reported that both methods yield extracellular vesicles in the size range of exosomes, which can be used in downstream analyses. Nanoparticle Tracking Analysis (NTA) is used for measuring exosome particles concentration, and WB or ELISA can be used in exosomal biomarkers analysis. Creative Biostructure standard exosome products guarantee higher purity and quality to meet our customer research.</p>
<b>Description</b>	Animal plasma exosome, Animal derived EV
<b>Form</b>	Lyophilized powder/ frozen liquid
<b>Concentration</b>	>1x10 <sup>6</sup> particles
<b>Storage</b>	Store at -20°C or colder. Recommend to avoid repeated freeze-and-thaw cycles.
<b>Reconstitution</b>	Reconstitute lyophilized exosome by adding deionized water for a desired final concentration. Centrifuge before opening to ensure exosomes are at the bottom, resuspend exosomes by pipetting and/or vortex, please avoid bubbles. Centrifuge again and mix well for use.