



## HQExo<sup>TM</sup> Microvesicles-NCI-H1975

## **Catalog: LEV-14**

## **PRODUCT INFORMATION**

| Name             | HQExo <sup>™</sup> Microvesicles-NCI-H1975   |
|------------------|--|
| Cat No.          | LEV-14   |
| Source           | Microvesicles derived from human lung carcinoma (non-small cells) (NCI-H975 cell line)   |
| Product Overview | Microvesicles are a type of extracellular vesicles (EVs) that are derived by cell membrane blebbing with a dia<br>meter from 100 nm to 1000 nm. While exosomes are smaller with a diameter between 30-160 nm and released<br>by cell exocytosis. Microvesicles involve in intercellular cross-talk and can transport molecules such as mRN<br>A, miRNA, lipids and proteins between cells, which make microvesicle play an important role in disease diagn<br>osis. Due to its molecular transfer function, circulating microvesicles may be useful for the delivery of drugs to<br>specific target cells. HQExo <sup>™</sup> microvesicles isolated from cancer cell lines could use as positive controls for E<br>LISA, FACS, WB. It has been reported that microvesicle express CD40, selectins, integrins, and cytoskeletal p<br>roteins, and their membranes are highly enriched in cholesterol, phosphatidylserine, and diacylglycerol. Micro<br>vesicles/exosomes has attracted more and more attention to anti-cancer research and regeneration. Microvesicl<br>es can be purified by ultracentrifugation and precipitation, then characterized by nanoparticles tracking analysi<br>s (NTA) and ELISA or WB. Lyophilization is useful for a long-term storage at 4°C, and frozen liquid should b<br>e kept at -20°C to -80°C. Creative Biostructure standard microvesicles products guarantee higher purity and qu<br>ality to meet our customer's downstream analyses. |
| Form             | Lyophilized powder/ frozen liquid  |
| Concentration    | >1x10^6 particles  |
| Storage          | Lyophilized powder store at 4 °C. Resuspension store at -80°C. Recommended to avoid repeated freeze-and-th raw cycles.   |