



Lyophilized DOPC/DOPE/CHOL ATP Liposome, NB D-lipid Labeled

Catalog: Lipo-216RG

PRODUCT INFORMATION

Name	Lyophilized DOPC/DOPE/CHOL ATP Liposome, NBD-lipid Labeled
Cat No.	Lipo-216RG
Product Overview	The encapsulation of ATP in liposomes markedly promotes its effectiveness by preventing the hydrolysis by extracellular enzymes, increasing ATP circulation time and enhancing its intracellular penetration. ATP liposomes can be used in various models such as myocardial, liver, retina and wound healing ischemia. Studies have shown the ability of liposomal encapsulated ATP to prevent cell death and tissue dysfunction following ischemic events. The concentration of encapsulated ATP is 0.5µmol/vial. Creative Biostructure could customize different DOPC/DOPE/CHOL ratio to meet your requirements. We can also manufacture empty lyophilized liposomes (without ATP) for control with the same lipid composition as your desired.
Lipid Composition	DOPC/DOPE/CHOL/NBD-PE (0.735/0.45/0.30/0.015 µmol/vial) DOPC: 1,2-dioleoyl-sn-glycero-3-phosphoc holine DOPE: 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine CHOL: Cholesterol NBD PE: 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-(7-nitro-2-1,3-benzoxadiazol-4-yl) (ammonium salt) (NBD PE)
Form	Lyophilized Powder
Storage Buffer	PBS, pH 7.4 with trehalose as lyoprotectant
Concentration	Lipid Concentration 1.5 µmol/vial
Stability	6 months
Storage	-20°C