



# MARK1 Protein Crystal


Catalog: CBCRY40

## PRODUCT INFORMATION

Name	MARK1 Protein Crystal
Cat No.	CBCRY40
Fragment	Residues 38-364
Protein Description	MAP/microtubule Affinity-regulating Kinase 1
Background	<p>The microtubule-associated protein (MAP)/microtubule affinity regulating kinase (MARK)/Par-1 phosphorylates microtubule-associated proteins tau, MAP2, and MAP4 and is involved in the regulation of microtubule-based transport. There are four isoforms of MARK in the human kinome that form a subfamily of the Snf1/AMP-activated protein kinase family of kinases within the calcium/calmodulin-dependent protein kinase group. MARK kinases are relatively large; the longest isoform, MARK1, comprises 795 amino acids. It is involved in the regulation of neuronal migration through its dual activities in regulating cellular polarity and microtubule dynamics, possibly by phosphorylating and regulating DCX. MARK1 also acts as a positive regulator of the Wnt signaling pathway, probably by mediating phosphorylation of dishevelled proteins (DVL1, DVL2 and/or DVL3).</p>
Protein Classification	Signaling Protein, Transferase
Structure Weight	301702.41 Da
Method	X-Ray Diffraction
Resolution	2.60 Å
Reference	<p>Marx A, Nugoor C, Mueller J, Panneerselvam S, Timm T, Bilang M, Mylonas E, Svergun DI, Mandelkow E-, Mandelkow EJ.Biol.Chem. Structural variations in the catalytic and ubiquitin-associated domains of microtubule-associated protein/microtubule affinity regulating kinase (Mark) 1 and mark2. (2006) 281 p.27586</p>

## USAGE GUIDELINES

General	Avoid excessive mixing or shocking to prevent aggregation. Long term storage above -80°C may result in aggregate formation.
---------	---



<b>Storage</b>	Short term: +2°C to +8°C
	Long term: -80°C
<b>Stability</b>	n.a.
<b>Freezing</b>	Can be frozen, but avoid multiple freeze/thaw cycles.